HEALTHY LIVER WITH BETTER DIGESTION

HEPASEL

INJECTION













HEALTHY LIVER WITH BETTER DIGESTION

HEPASEL

Injection

Composition:

Each ml contains Phenoxy - 2 Methyl - 2 Propionic Acid 100 mg



Cattle, Sheep & Goat: Hepatic Failure, Indigestion, Acetonemia, In-appetence, Bloat, Alimentary toxicosis, Rumen impaction, Dyspepsia with Meteorism, Ketosis, as an adjuvant therapy during treatment of liver fluke disease.

Horses: Liver disorders due to imbalanced feed, hepatic failure due to Piroplasmosis and Leptospirosis.

Dogs: Icterus, hepatic failure, as an adjuvant therapy in the treatment of Leptospirosis and Canine Distemper.

Pharmacological Properties

After parenteral administration Hepasel injection quickly absorbed in the blood stream and characterized by an

elective action on liver, with an increase in bile secretion and favoring the performance of digestive functions. Hepasel injection performs its secretory stimulatory action by acting directly on the glandular system without excitation or depression of central nervous system (CNS) or autonomous nervous system. Hepasel injection also have choleretic activity by stimulating bile production by the liver, therefor indicated in the diseases associated with hepatic insufficiency. Hepasel injection is excreted unmodified through urine and feces.

Dosage & Administration:

Administer the following dose by deep intramuscular, intraperitoneal route or slow intra-venous route. The dose can be repeated after every 24 hours according to veterinarian advice.

Cattle, Horses: 10 ml per 100 kg body weight. Sheep, Goat & Dogs: 1 ml per 10 kg bodyweight.

Precautions:

Do not mix with solutions containing calcium salts. Store below 25°C in a cool and dry place.

Packing: 50 ml.

بیک وقت کیلٹیم کے ساتھ استعال نہ کریں۔ 25°C سے کم ٹھنڈی اور خشک جگہ پر نہ رکھیں۔





علامات : هیپاسیل انجیکش جگر،معده، پیة اورلبلبه ک افعال کودرست کرتا ہے۔ قبض، بھوک میں کمی، بدہضمی، ابچارہ، ٹمپنی، بلوٹ، امپیکشن، کیٹوسیز، وائرس کی بیاریوں سے جگر پر ہونے والےمضراثرات کوزائل کرنے کے لئے ھیپاسیل کو علاج اور معاون علاج کے طور پر استعال کیا جاسکتا ہے۔

اجزاء؛ برملي ليثر ميں شامل ہيں فينوكس - 2 ميتهائل - 2 يرويائيونك ايسدُّ 100 ملى كرام خوراك وطريقه استعال: درج ذیل خوراک پیٹوں یاورید میں استعال کریں۔

گائے ، بھینس ، گھوڑے : 10 ملی لیٹر فی 100 کلوگرام وزن۔ بھیٹر، بکری، کتے: 1 ملی لیٹر فی 10 کلوگرام وزن۔

علامات کومد نظرر کھتے ہوئے ڈاکٹر کی ہدایات کے مطابق خیں کے مصرف

خوراک ہر 24 گھنٹے بعد دوبارہ استعمال کریں۔





