IZOVAC B1 HITCHNER

FREEZE-DRIED LIVE VACCINE AGAINST NEWCASTLE DISEASE

COMPOSITION :

Each dose of vaccine contains: live attenuated Newcastle Disease virus, strain B1 Hitchner: 10^{6.5} EID₅₀ in stabilizing solution.

INDICATIONS :

Active immunization against Newcastle Disease.

DOSAGE AND ADMINISTRATION :

Administer by eye drop, spray or via drinking water. First administration: 1 day of age (spray or eye drop). Second administration: 2-3 weeks later. Booster: every 2-3 months in layers or breeders. The amount of water changes according to the age or season.

WARNINGS :

Via spray:

- utilize a sprayer producing droplets of about 50 micron
- spray at about 1 meter of distance from the birds

Via drinking water :

- Suspend 2-3 hours before vaccination the administration of drinking water, in order to make the animals partially thirsty
- Control that the troughs are empty, clean and in particular that they do not contain residues of poultry feed, faeces or traces of disinfectants
- After determining the number of birds to be vaccinated prepare in clean containers (without disinfectans) the quantity of water to be drunk necessary for the administration of the vaccine
- All the birds should have the access to drinkers at the same time
- The reconstitution of the vaccine, the final dilution in the drinking water and the administration must be effected in the shortest time
- # The drinking water containing the vaccine must be exhausted in a period ranging from 2 to 4 hours
- Do not expose the water containing the vaccine to heat sources (incubating lamps, sun and so on) or to intense light
- # Use water not containing chlorine or traces of disinfectans
- # Skim milk powder may be added to the water (2-3 grams per litre)

Via eye drop :

Administer one drop of vaccine suspension in to one eye by means of the dropper.

STORAGE : Store at +2/+8°C.

EXPIRY TIME : 2 years.

PACKING : Box containing 10 vials of 1000 doses.



For Veterinary Use Only

