

IZOVAC CORYZA 3

EMULSIFIED INACTIVATED VACCINE AGAINST INFECTIOUS CORYZA

COMPOSITION :

Each dose of 0.5 ml of vaccine contains Inactivated cultures of :

Haemophilus paragallinarum serotype A : 5×10^9 CFU;

Haemophilus paragallinarum serotype B : 5×10^9 CFU;

Haemophilus paragallinarum serotype C : 5×10^9 CFU; in oil emulsion

INDICATIONS :

Active immunization of breeders and layers against Infectious Coryza.

DOSAGE AND ADMINISTRATION :

Inject subcutaneously (at the back of the neck) or intramuscularly (pectoral muscles) of breeders and laying hens. First vaccination at 6-8 weeks, booster dose 4 weeks before point of lay. The second vaccination can be done alone or in association with other antigens.

WARNINGS :

- ☛ Let the vaccine to reach the room temperature (+20°C) before use
- ☛ Shake well before and during use
- ☛ Only healthy birds should be vaccinated
- ☛ Accidental injection of this vaccine into the vaccinator can cause serious localized reaction for which expert medical advice should be sought at once, informing the doctor that the vaccine is an oil emulsion

STORAGE :

Store at +2/+8°C. Do not freeze.

EXPIRY TIME :

2 years.

PACKING :

Box containing 10 bottles of 1000 doses.

For Veterinary Use Only

Marketed by:
Animal Health Division
RENATA LIMITED
Mirpur, Dhaka, Bangladesh



Manufactured by:
IZO S.r.l a socio unico
Sede legale: Via San Zeno, 99/A- 25124 Brescia, Italy