# **IZOVAC LASOTA**

### FREEZE-DRIED LIVE VACCINE AGAINST NEWCASTLE DISEASE

#### **COMPOSITION:**

Each dose of vaccine contains:

Live attenuated Newcastle Disease Virus, strain Lasota: 10<sup>6</sup> EID<sub>50</sub> in stabilizing solution.

#### **INDICATIONS:**

The vaccine is recommended for booster vaccination against Newcastle Disease.

## **DOSAGE AND ADMINISTRATION:**

Administer via drinking water. The vaccine is administered at 21-28 days of age and then every 2-3 months in layers or breeders. The vaccine Izovac B1 Hitchner or Izovac Clone must be administered at 1 day of age as first vaccination. The amount of water changes according to the age or season.

#### **WARNINGS:**

- ¶ Suspend 2-3 hours before vaccination the administration of drinking water, in order to make
  the birds partially thirsty
- ♠ Control that the troughs are empty, clean and in particular that they do not contain residues of poultry feed, faeces or traces of disinfectants
- ♠ After determining the number of birds to be vaccinated prepare in clean containers (without disinfectans) the quantity of water to be drunk necessary for the administration of the vaccine
- ♠ All the birds should have the access to drinkers at the same time
- ₱ The reconstitution of the vaccine, the final dilution in the drinking water and the administration must be effected in the shortest time
- ₱ The drinking water containing the vaccine must be exhausted in a period ranging from 2 to 4 hours.
- Do not expose the water containing the vaccine to heat sources (incubating lamps, sun and so on) or to intense light
- Use water not containing chlorine/traces or disinfectans
- ¶ Skim milk powder may be added to the water (2-3 grams per litre)

**STORAGE**: Store at +2/+8°C.

**EXPIRY TIME**: 2 years.

PACKING:

Box containing 10 vials of 1000 doses.

For Veterinary Use Only



