IZOVAC ND - EDS

EMULSIFIED INACTIVATED VACCINE AGAINST NEWCASTLE DISEASE AND EDS '76

COMPOSITION:

Each dose of 0.5 ml of vaccine contains: Inactivated Newcastle Disease Virus: 50 PD₅₀;

Inactivated EDS '76 Virus: 108 EID50 before inactivation; in oil emulsion

INDICATIONS:

Active immunization against Newcastle Disease and Egg Drop Syndrome (EDS '76).

DOSAGE AND ADMINISTRATION:

Inject subcutaneously (at the back of the neck) or intramuscularly (pectoral muscles) at 18-20 weeks of age before the laying period. A single administration is sufficient to protect the birds against EDS, while in the case of Newcastle Disease, the administration should be preceded by other vaccinations with live or inactivated vaccine, as is usual in the prophylaxis of this infection.

WARNINGS:

- ♠ Let the vaccine reach the room temperature (+20°C) before use
- ¶ Shake well before and during use
- ₱ Only healthy birds should be vaccinated
- ♠ Accidental injection of this vaccine into the vaccinator can cause serious localized reaction
 for which expert medical advice should be sought at once, informing the doctor that the
 vaccine is an oil emulsion

STORAGE:

Store at +2/+8°C. Do not freeze.

EXPIRY TIME:

2 years.

PACKING:

Box containing 10 bottles of 1000 doses.

For Veterinary Use Only



