

SUPRAVAC 10

Inactivated vaccine against
Respiratory & Reproductive Diseases



SUPRAVAC 10

Inactivated vaccine against
Respiratory & Reproductive Diseases



Composition

Quali-quantitative Formula	
Infectious Bovine Rhinotracheitis Virus	$10^{3,4}$ TCID ₅₀
Bovine Parainfluenza Virus Type 3	$10^{3,4}$ TCID ₅₀
Bovine Respiratory Syncytial virus	10^3 TCID ₅₀
Inactivated Cytopathic sample Bovine Viral Diarrhea	10^3 TCID ₅₀
Inactivated Cytopathic sample Bovine Viral Diarrhea Type 2	10^3 TCID ₅₀
Inactivated Non-Cytopathic sample Bovine Viral Diarrhea	10^3 FAID ₅₀
Bacterin of Leptospira canicola	10^6 germs
Bacterin of Leptospira icterohaemorrhagiae	10^6 germs
Bacterin of Leptospira pomona	10^6 germs
Bacterin of Leptospira grippothyphosa	10^6 germs
Bacterin of Leptospira hardjo	10^6 germs
Bacterin of Leptospira wolffii	10^6 germs
Bacterin of Pasteurella haemolytica	10^4 germs

Administration

Route: subcutaneously

Dosage

Dose: 5ml Adult Cattle & 3ml in Calves

Booster : after 30 days

Pregnant females should be vaccinated 30 days before parturition