



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

Elanco™

AF1375

Maxiban™

TM

Narasin/Nicarbazin

Anticoccidial Premix

ACTIVE CONSTITUENTS: NARASIN 80 g/kg
NICARBAZIN 80 g/kg

For the prevention of coccidiosis caused by
Eimeria acervulina, *E. brunetti*, *E. maxima*,
E. mivati, *E. necatrix*, and *E. tenella*.

FOR USE IN BROILER CHICKEN FEEDS ONLY.
25 kg NET

DISPOSAL OF BAGS: Shake and empty contents into medicated feed.
Do not dispose of undiluted chemicals on site. Puncture or shred and bury
empty bags in a local authority landfill. If not available bury the bag below
500 mm in a disposal pit specifically marked and set up for this purpose
clear of waterways, vegetation and roots. Empty bags and product should
not be burned.

Store below 30°C (Room Temperature) in a dry place.
For expiry date and batch number see side of bag

APVMA Approval No. 36802/25kg/0610

ELANCO ANIMAL HEALTH
A Division of Eli Lilly Australia Pty Limited
A.B.N. 39 000 233 992
112 WHARF ROAD, WEST RYDE, N.S.W. 2114
TEL: Toll Free 1800 226 324



USE NO HOOKS

BG090038ALX

DIRECTIONS FOR USE:

Restraints:

EGGS: DO NOT USE in birds which are producing or may in the
future produce eggs or egg products for human consumption.

Contra-indications:

DO NOT USE in turkeys or game birds – may be fatal.
DO NOT USE in horses or other equines – may be fatal.
Poultry consuming narasin should not be treated with tiamulin.
Severe growth depression or death may occur.
DO NOT USE concurrently with another ionophore, as severe growth
depression or death may occur.

Precautions:

Nicarbazin medicated broilers may show reduced heat tolerance
if exposed to high temperature and high humidity.
Provide adequate drinking water and ventilation during these periods.

MIXING DIRECTIONS:

**IMPORTANT – THIS PREMIX MUST BE THOROUGHLY MIXED
IN FEED BEFORE USE.**

Thoroughly mix 0.5 to 0.625 kg MAXIBAN Premix in one tonne of feed
to provide 40 to 50 ppm of each active ingredient per tonne of complete
feed.

FEEDING DIRECTIONS:

Feed continuously as the only ration for at least 14 days.

WITHHOLDING PERIOD:

MEAT: NIL

EGGS: DO NOT USE in birds which are producing or may in
the future produce eggs or egg products for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI):

This product does not have an ESI established.
For advice on the ESI, contact Elanco Animal Health (Australia)
on 1800 226 324.

SAFETY DIRECTIONS:

Poisonous if swallowed. Will damage eyes and will irritate the nose,
throat and skin. Avoid contact with eyes and skin. Do not inhale
dust when opening the container and mixing into feed. Repeated
exposure may cause allergic disorders. Wear cotton overalls
buttoned to the neck and wrist and a washable hat, elbow-length
PVC gloves, goggles and disposable dust mask covering mouth and
nose. After each day's use, wash gloves, goggles and contaminated
clothing. After use and before eating, drinking or smoking, wash
hands, arms and face thoroughly with soap and water.

**FIRST AID: If poisoning occurs, contact a doctor or Poisons
Information Centre. Phone Australia 13 1126. If in eyes, hold eyes
open, flood with water for at least 15 minutes and see a doctor.**

Additional information is in the Safety Data Sheet.

Elanco, Maxiban and the diagonal bar are trademarks owned or
licensed by Eli Lilly and Company, its subsidiaries or affiliates.

**May cause damage to organs (Heart, Nervous system,
Muscle) through prolonged or repeated exposure.**
IF SWALLOWED: Rinse mouth. IF ON SKIN: Wash with plenty of
soap and water. If skin irritation occurs: Get medical advice/attention.
IF INHALED: If breathing is difficult, remove victim to fresh air and
keep at rest in a position comfortable for breathing. If experiencing
respiratory symptoms: Call a POISON CENTRE/doctor.

BG090038ALX

**Elanco™
Maxiban™**

**Elanco™
Maxiban™**

**Elanco™
Maxiban™**