BRONGOWEEX Prescribing Information POWDER

ACTION

BRONCOVEEX is a soluble powder for oral administration through drinking water for the treatment of respiratory diseases with thickened bronchial secretions. Being mucolytic, it fluidifies the mucus and improves the state of the respiratory tract, enabling the penetration and diffusion of the curing agents.

INDICATIONS

BRONCOVEEX is used in cattle, horses, sheep, goats and poultry for treatment in respiratory diseases, such as:

Pharyngitis, Laryngitis, Bronchitis, Tracheitis,

Pleuritis, Tracheobronchitis, Pneumonia, Broncho-

Pneumonia, Bronchiectasis, Atelectasis, Mumps,

Lung abscesses, CRD & Rhinitis.

In acute infections, BRONCOVEEX is recommended to be given with antibiotics, antibacterials, etc.

COMPOSITION

Each 100 gm contains:

Bromhexine Hydrochlorate

1gm

Tartaric acid

15 gm

DOSAGE AND ADMINISTRATION

BRONCOVEEX is administered orally through feed or drinking water as follows:

Cattle

4-8 gm/100kg live weight

Horses

4-8 gm/100 kg live weight

Young cattle, sheep & goats

10-20 gm/100 kg live weight

Poultry

1 gm/litre of drinking water per day.

BRONCOVEEX should be continued for 3 to 5 days accompanied by treatment based on antibiotics, antibacterials, etc.

SIDE-EFFECTS

None reported on therapeutic doses.

CONTRA-INDICATIONS

BRONCOVEEX should not be given to animals with ulcerous processes in the digestive tract. Solution of sodium chloride should also not be administered with BRONCOVEEX.

CAUTIONS

Treatment should be stopped 3 days before animals are to be slaughtered. Suppression of treatment is not necessary in milk producing animals or egg producing poultry.

PACKS

BRONCOVEEX is available in packs of 100gm and one kg.



