

Time tested broad spectrum antibacterial

TRYTON

(Trimethoprim, Sulphadiazine) Oral Suspension

An effective combination to chick bacterial growth.

An effective combination to limit diseases / common poultry diseases.

An effective combination for maximum profit through better health.

Combined effective protectional formula from day one.

Protection from day one.

An effective combination for better health.

An effective combination for common poultry diseases.

Higher gain with proven results

Coli Septicemia,
Infectious Coryza,
Fowl Typhoid,
Fowl Cholera and
Staphylococal arthritis












Formulation

Trimethoprim 80mg/ml Sulphadiazine 400mg/ml

- ☐ Time tested broad spectrum antibacterial for the treatment of bacterial disease in poultry specially coli septicemia, infectious coryza, Fowl Typhoid, Fowl cholera and staphylococcal arthritis.
- ☐ It is rapidly absorbed and widely distributed.
- ☐ Has Unique mode of action

(Double sequential blockade)



- ☐ Due to this double sequential blockade chance of resistance very low.
- ☐ Has micronized particle easily dispersible and uniform mixing.
- Does not affect egg production and growth rate.

Dosage and Administration

- ☐ 15 30 mg per kilogram of body weight
- ☐ 1ml/4.5liters of water for 5 days
- ☐ Use 5ml of Tryton 13-22 liters of drinking water for 1000 chicks daily. This treatment should continue for 5 days.

Precaution

Pack

Shake well before use.

■ Available in 100ml, 250ml and 1 Litre bottles.

